

2000**Prolonged exposure to intermittent alcohol vapors blunts hypothalamic responsiveness to immune and non-immune signals.**

S.Lee, D.Schmidt, F.Tilders, M.Cole, A.Smith and C.River,
Alcoholism: Clinical and Experimental Research 24 (1) 110-122 (2000)

Impaired protein synthesis induced by acute alcohol intoxication is associated with changes in eIF4E in muscle and eIF2B in liver.

C.H.Lang, R.A.Frost, V.Kumar, D.Wu and T.C.Vary,
Alcoholism: Clinical and Experimental Research, 24 (3) 322-331 (2000)

Impairment in eyeblink classical conditioning in adult rats exposed to ethanol as neonates.

J.T.Green, R.F.Rogers, C.R.Goodlett and J.E.Steinmetz,
Alcoholism: Clinical and Experimental Research, 24 (4) 438-447 (2000)

2001**Mice that lack corticotropin-releasing factor (CRF) receptors Type 1 show a blunted ACTH response to acute alcohol despite up-regulated constitutive hypothalamic CRF gene expression.**

S. Lee, G.W. Smith, W. Vale, K-F. Lee and C. Rivier,
Alcoholism: Clinical and Experimental Research, 25 (3) 427-433 (2001)

Local cerebral glucose utilization rates in alcohol-naïve High-Alcohol-Drinking (HAD) and Low-Alcohol-Drinking (LAD) rats.

J.E.Learn, D.G.Smith, W.J.McBride, L.Lumeng and T-K.Lee,
Alcoholism: Clinical and Experimental Research, 25 (4) 517-523 (2001)

Responsivity and development of tolerance to the motor impairing effects of moderate doses of ethanol in alcohol-preferring (P) and –nonpreferring (NP) rat lines

R.E.Bell et al,
Alcoholism: Clinical and Experimental Research, 25 (5) 644-650 (2001)

MK-801 can exacerbate or attenuate behavioral alterations associated with neonatal alcohol exposure in the rat, depending on the timing of administration

J.D.Thomas, S.L.Fleming and E.P.Riley,
Alcoholism: Clinical and Experimental Research, 25 (5) 764-773 (2001)

Ethanol-induced impairments in spatial working memory are not due to deficits in learning

S.E.Hoffmann and D.B.Matthews,
Alcoholism: Clinical and Experimental Research, 25 (6) 856-861 (2001)

Effects of concurrent access to multiple ethanol concentrations and repeated deprivations on alcohol intake of alcohol-preferring rats

Z.A.Rodd-Henricks et al,
Alcoholism: Clinical and Experimental Research, 25 (8) 1140-1150 (2001)

Early alteration in leukocyte populations and Th1/Th2 function in ethanol-consuming mice

S.Starkenbergh, M.E.Munroe and C.Waltenbaugh,
Alcoholism: Clinical and Experimental Research, 25 (8) 1221-1230 (2001)

Differential response to the aversive properties of alcohol in alcohol-preferring (sP) and –nonpreferring (sNP) rats

G.Brunetti, S.Mellis, S.Serra, G.Vacca, G.Colombo and G.L.Gessa, Poster presentation, 8th Congress of the European Society for Biomedical Research on Alcoholism, September (2001)

Increased ethanol self-administration in δ -opioid receptor knockout mice

A.J.Roberts et al,
Alcoholism: Clinical and Experimental Research, 25 (9) 1249-1256 (2001)

2001**Alcohol-Naïve alcohol-preferring (P) rats exhibit higher local cerebral glucose utilization than alcohol-nonpreferring (NP) and Wistar rats**

D.G.Smith, J.E.Learn, W.J.McBride, L.Lumeng, T-K.Li and J.M.Murphy,
Alcoholism: Clinical and Experimental Research, 25 (9) 1309-1316 (2001)

Evidence that the Lore-1 region specifies ethanol-induced activation in addition to sedative/hypnotic sensitivity to ethanol

I.C.Owens, B.Bennett and T.E.Johnson,
Alcoholism: Clinical and Experimental Research, 25 (11) 1551-1557 (2001)

2002**Selective and enduring deficits in spatial learning after limited neonatal binge alcohol exposure in male rats**

T.B.Johnson and C.R.Goodlett,
Alcoholism: Clinical and Experimental Research, 26 (1) 83-93 (2002)

Reciprocal congenics defining individual quantitative trait loci for sedative/hypnotic sensitivity to ethanol

B.Bennett, M.Beeson, L.Gordon and T.E.Johnson,
Alcoholism: Clinical and Experimental Research, 26 (2) 149-157 (2002)

Altered EEG responses to ethanol in adult rats exposed to ethanol during adolescence

C.J.Slawecki,
Alcoholism: Clinical and Experimental Research, 26 (2) 246-254 (2002)

Microdialysis of dopamine in the nucleus accumbens of alcohol-preferring(P) rats during anticipation and operant self-administration of ethanol

R.I. Melendez, Z.A. Rodd-Henricks, E.A. Engleman, T-K Li, W.J. McBride and J.M. Murphy, Alcoholism: Clinical and Experimental Research, 26 (3) 318-325 (2002)

Chronic ethanol treatment and withdrawal alter ACPD-evoked calcium signals in developing Purkinje neurons

J.G. Netzeband, J.R. Schneeloch, C. Trotter, J.N. Caguioa-Aquino and D.L. Gruol, Alcoholism: Clinical and Experimental Research, 26 (3) 386-393 (2002)

Binge ethanol exposure in adult rats causes necrotic cell death

J.A. Obernier, T.W. Bouldin and F.T. Crews,
Alcoholism: Clinical and Experimental Research, 26 (4) 547-557 (2002)

The decreased cellular expression of neuropeptide protein in rat brain structures during ethanol withdrawal after chronic ethanol exposure

Roy and S.C. Pandey,
Alcoholism: Clinical and Experimental Research, 26 (6) 796-803 (2002)

Chronic-intermittent ethanol exposure during adolescence prevents normal developmental changes in sensitivity to ethanol-induced motor impairments

A.M. White, J.G. Bae, M.C. Truesdale, S. Ahmad, W.A. Wilson and H.S. Swartwelder, Alcoholism: Clinical and Experimental Research, 26 (7) 960-968 (2002)

Possible pleiotropic effects of genes specifying sedative/hypnotic sensitivity to ethanol on other alcohol-related traits

J.C. Owens, B. Bennett and T.E. Johnson,
Alcoholism: Clinical and Experimental Research, 26 (10) 1461-1467 (2002)

2003

The neurotoxicity induced by ethanol withdrawal in mature organotypic hippocampal slices might involve cross-talk between metabotropic glutamate Type 5 receptors and N-methyl-D-aspartate receptors.

B.R. Harris et al,

Alcoholism: Clinical and Experimental Research, 27 (11) 1724-1735 (2003)

Comparison between the effect of systematic and intracerebroventricular injection of this drug on pituitary and hypothalamic responses

S. Lee, D. Selvage, K. Hansen and C. Rivier,

Endocrinology, 145 (10) 4470-4479 (2004)

2005

Sex differences in ethanol liquid diet consumption in Sprague-Dawley rats

M.R. Piano, T.M. Carrigan and D.W. Schwertz,

Alcohol, 35 113-118 (2005)

2008

Neuropeptide y in the central nucleus of the amygdala suppresses dependence-induced increases in alcohol drinking

N. W. Gilpin, Kaushik Misra, and George F. Koob

Pharmacol Biochem Behav. 90(3), 475 -480 (2008)

2009

Intra-cornu ammonis 1 administration of the human immunodeficiency virus-1 protein trans-activator of transcription exacerbates the ethanol withdrawal syndrome in rodents and activates N-methyl-D-aspartate glutamate receptors to produce persisting spatial learning deficits

R.L. Self, K.J. Smith, T.R. Butler, J.R. Pauly and M.A. Prendergast

Neuroscience, 163, 868-876, (2009)

Permanent impairment of birth and survival of cortical and hippocampal proliferating cells following excessive drinking during alcohol dependence

H. N. Richardson, S. H. Chan, E. F. Crawford, Y.K. Lee, C. K. Funk, G. F. Koob, and C. D. Mandyam Neurobiol Dis. 36 (1), 1-10, (2009).