

## 2000

**Prolonged exposure to intermittent alcohol vapors blunts hypothalamic responsiveness to immune and non-immune signals.**

S.Lee, D.Schmidt, F.Tilders, M.Cole, A.Smith and C.River,  
Alcoholism: Clinical and Experimental Research 24 (1) 110-122 (2000)

**Impaired protein synthesis induced by acute alcohol intoxication is associated with changes in eIF4E in muscle and eIF2B in liver.**

C.H.Lang, R.A.Frost, V.Kumar, D.Wu and T.C.Vary,  
Alcoholism: Clinical and Experimental Research, 24 (3) 322-331 (2000)

**Impairment in eyeblink classical conditioning in adult rats exposed to ethanol as neonates.**

J.T.Green, R.F.Rogers, C.R.Goodlett and J.E.Steinmetz,  
Alcoholism: Clinical and Experimental Research, 24 (4) 438-447 (2000)

## 2001

**Mice that lack corticotropin-releasing factor (CRF) receptors Type 1 show a blunted ACTH response to acute alcohol despite up-regulated constitutive hypothalamic CRF gene expression.**

S. Lee, G.W. Smith, W. Vale, K-F. Lee and C. Rivier,  
Alcoholism: Clinical and Experimental Research, 25 (3) 427-433 (2001)

**Local cerebral glucose utilization rates in alcohol-naïve High-Alcohol-Drinking (HAD) and Low-Alcohol-Drinking (LAD) rats.**

J.E.Learn, D.G.Smith, W.J.McBride, L.Lumeng and T-K.Lee,  
Alcoholism: Clinical and Experimental Research, 25 (4) 517-523 (2001)

**Responsivity and development of tolerance to the motor impairing effects of moderate doses of ethanol in alcohol-preferring (P) and –nonpreferring (NP) rat lines**

R.E.Bell et al,  
Alcoholism: Clinical and Experimental Research, 25 (5) 644-650 (2001)

**MK-801 can exacerbate or attenuate behavioral alterations associated with neonatal alcohol exposure in the rat, depending on the timing of administration**

J.D.Thomas, S.L.Fleming and E.P.Riley,  
Alcoholism: Clinical and Experimental Research, 25 (5) 764-773 (2001)

**Ethanol-induced impairments in spatial working memory are not due to deficits in learning**

S.E.Hoffmann and D.B.Matthews,  
Alcoholism: Clinical and Experimental Research, 25 (6) 856-861 (2001)

**Effects of concurrent access to multiple ethanol concentrations and repeated deprivations on alcohol intake of alcohol-preferring rats**

Z.A.Rodd-Henricks et al,  
Alcoholism: Clinical and Experimental Research, 25 (8) 1140-1150 (2001)

**Early alteration in leukocyte populations and Th1/Th2 function in ethanol-consuming mice**

S.Starkenbergh, M.E.Munroe and C.Waltenbaugh,  
Alcoholism: Clinical and Experimental Research, 25 (8) 1221-1230 (2001)

**Differential response to the aversive properties of alcohol in alcohol-preferring (sP) and –nonpreferring (sNP) rats**

G.Brunetti, S.Mellis, S.Serra, G.Vacca, G.Colombo and G.L.Gessa, Poster presentation, 8th Congress of the European Society for Biomedical Research on Alcoholism, September (2001)

**Increased ethanol self-administration in  $\delta$ -opioid receptor knockout mice**

A.J.Roberts et al,  
Alcoholism: Clinical and Experimental Research, 25 (9) 1249-1256 (2001)

**2001****Alcohol-Naïve alcohol-preferring (P) rats exhibit higher local cerebral glucose utilization than alcohol-nonpreferring (NP) and Wistar rats**

D.G.Smith, J.E.Learn, W.J.McBride, L.Lumeng, T-K.Li and J.M.Murphy,  
Alcoholism: Clinical and Experimental Research, 25 (9) 1309-1316 (2001)

**Evidence that the Lore-1 region specifies ethanol-induced activation in addition to sedative/hypnotic sensitivity to ethanol**

I.C.Owens, B.Bennett and T.E.Johnson,  
Alcoholism: Clinical and Experimental Research, 25 (11) 1551-1557 (2001)

**2002****Selective and enduring deficits in spatial learning after limited neonatal binge alcohol exposure in male rats**

T.B.Johnson and C.R.Goodlett,  
Alcoholism: Clinical and Experimental Research, 26 (1) 83-93 (2002)

**Reciprocal congenics defining individual quantitative trait loci for sedative/hypnotic sensitivity to ethanol**

B.Bennett, M.Beeson, L.Gordon and T.E.Johnson,  
Alcoholism: Clinical and Experimental Research, 26 (2) 149-157 (2002)

**Altered EEG responses to ethanol in adult rats exposed to ethanol during adolescence**

C.J.Slawecki,  
Alcoholism: Clinical and Experimental Research, 26 (2) 246-254 (2002)

**Microdialysis of dopamine in the nucleus accumbens of alcohol-preferring(P) rats during anticipation and operant self-administration of ethanol**

R.I. Melendez, Z.A. Rodd-Henricks, E.A. Engleman, T-K Li, W.J. McBride and J.M. Murphy, Alcoholism: Clinical and Experimental Research, 26 (3) 318-325 (2002)

**Chronic ethanol treatment and withdrawal alter ACPD-evoked calcium signals in developing Purkinje neurons**

J.G. Netzeband, J.R. Schneeloch, C. Trotter, J.N. Caguaioa-Aquino and D.L. Gruol, Alcoholism: Clinical and Experimental Research, 26 (3) 386-393 (2002)

**Binge ethanol exposure in adult rats causes necrotic cell death**

J.A. Obernier, T.W. Bouldin and F.T. Crews,  
Alcoholism: Clinical and Experimental Research, 26 (4) 547-557 (2002)

**The decreased cellular expression of neuropeptide protein in rat brain structures during ethanol withdrawal after chronic ethanol exposure**

Roy and S.C. Pandey,  
Alcoholism: Clinical and Experimental Research, 26 (6) 796-803 (2002)

**Chronic-intermittent ethanol exposure during adolescence prevents normal developmental changes in sensitivity to ethanol-induced motor impairments**

A.M. White, J.G. Bae, M.C. Truesdale, S. Ahmad, W.A. Wilson and H.S. Swartwelder, Alcoholism: Clinical and Experimental Research, 26 (7) 960-968 (2002)

**Possible pleiotropic effects of genes specifying sedative/hypnotic sensitivity to ethanol on other alcohol-related traits**

J.C. Owens, B. Bennett and T.E. Johnson,  
Alcoholism: Clinical and Experimental Research, 26 (10) 1461-1467 (2002)

**2003**

**The neurotoxicity induced by ethanol withdrawal in mature organotypic hippocampal slices might involve cross-talk between metabotropic glutamate Type 5 receptors and N-methyl-D-aspartate receptors.**

B.R. Harris et al,

Alcoholism: Clinical and Experimental Research, 27 (11) 1724-1735 (2003)

**Comparison between the effect of systematic and intracerebroventricular injection of this drug on pituitary and hypothalamic responses**

S. Lee, D. Selvage, K. Hansen and C. Rivier,

Endocrinology, 145 (10) 4470-4479 (2004)

**2005**

**Sex differences in ethanol liquid diet consumption in Sprague-Dawley rats**

M.R. Piano, T.M. Carrigan and D.W. Schwertz,

Alcohol, 35 113-118 (2005)

**2008**

**Neuropeptide y in the central nucleus of the amygdala suppresses dependence-induced increases in alcohol drinking**

N. W. Gilpin, Kaushik Misra, and George F. Koob

Pharmacol Biochem Behav. 90(3), 475 -480 (2008)

**2009**

**Intra-cornu ammonis 1 administration of the human immunodeficiency virus-1 protein trans-activator of transcription exacerbates the ethanol withdrawal syndrome in rodents and activates N-methyl-D-aspartate glutamate receptors to produce persisting spatial learning deficits**

R.L. Self, K.J. Smith, T.R. Butler, J.R. Pauly and M.A. Prendergast

Neuroscience, 163, 868-876, (2009)

**Permanent impairment of birth and survival of cortical and hippocampal proliferating cells following excessive drinking during alcohol dependence**

H. N. Richardson, S. H. Chan, E. F. Crawford, Y.K. Lee, C. K. Funk, G. F. Koob, and C. D. Mandyam Neurobiol Dis. 36 (1), 1-10, (2009).